arXiv:1304.0164v1 [hep-ph] 31 Mar 2013

Generation of non-zero θ_{13} in broken A_4 neutrino mass model

Manikanta Borah,[*] Bikash Sharma,[†] and Mrinal Kumar Das[‡]

Department of Physics, Tezpur University, Tezpur-784028, India

Abstract

Stability of neutrino masses and mixings with non-zero θ_{13} is an important issue in recent neutrino physics. In context of this, a detail analysis on two-fold degenerate ($m_1 = m_2, m_3$) neutrino mass model is done with Renormaization Group Equation (RGE) in Minimal Supersymmetric Standard Model (MSSM). Considering the input values of neutrino masses and mixing angles at high scale predictions are studied at low energy scale using RGE. We have also found that two-fold degenerate model with $\theta_{13} = 0$ can produce consistent results at low energy with non-zero θ_{13} from the breaking of a generic A_4 model.

PACS numbers: 14.60.Pq, 11.10.Gh, 11.10.Hi

---

[*] mani@tezu.ernet.in

[†] bikash.sharma150@gmail.com

[‡] mkdas@tezu.ernet.in

I. INTRODUCTION

Results of recent neutrino oscillation experiments have provided concrete of evidence which confirms the existence of the non-zero yet tiny neutrino masses [1]. Recenet neutrinio oscillation experiments like T2K [2], Double ChooZ [3], Daya-Bay [4] and RENO [5] have not only made the earlier predictions for neutrino parameters more precise, but also predicted non-zero value of the reactor mixing angle θ_{13}. The latest global fit value for 3σ range of neutrino oscillation parameters [6] are as follows

$$\begin{aligned}
\Delta m^2_{21} &= (7.00 - 8.09) \times 10^{-5} \\
\Delta m^2_{31}(NH) &= (2.27 - 2.69) \times 10^{-3} \\
\Delta m^2_{23}(IH) &= (2.24 - 2.65) \times 10^{-3} \\
\sin^2\theta_{12} &= 0.27 - 0.34 \\
\sin^2\theta_{23} &= 0.34 - 0.67 \\
\sin^2\theta_{13} &= 0.016 - 0.030. \qquad (1)
\end{aligned}$$

The above recent data have positive evidence for non-zero θ_{13} as well, which was earlier thought to be zero or negligibly small. The value of this mixing θ_{13} has serious impact in recent neutrino physics as it leads to the non-zero CP violating phase in the neutrino sector. It has also non-trivial impact on neutrino mass hierarchy as studied in recent paper [7]. Detailed analysis of this non-zero θ_{13} using different approach is done viz. theoretical [8], phenomenological[9] before as well as after the discovery of this result in 2012. In view of the imortance of understanding the non-zero reactor mixing and CP violation in neutrino sector as mentioned above, present work is planned to present how a specific μ-τ symmetric mass matrix [10] with two-fold degenerate mass eigenvalues can produce non-zero θ_{13} along with Δm^2_{21} through radiative corrections. This work is also emphasised that how a broken A_4 symmetry at high scale can produce other neutrino oscillation parameters θ_{23}, θ_{12} and Δm^2_{23} at low energy scale through radiative corrections.

This article is organized as follows: in section II we discuss briefly the broken A_4 model which is used at high scale and about the RGE of mass eigen values and mixing to generate the non-zero reactor angle. In section III we discuss our numerical results and analysis. In the last section IV we outline the discussion and conclusions of our analysis.

II. NEUTRINO MASS MODELS AND RGE

A. A_4 model for neutrino mass

Type I seesaw framework is the simplest mechanism for generating tiny neutrino masses and mixing. The finite group of even permutation A_4 can explain the $\mu - \tau$ symmetric mass matrix obtained from this type I seesaw mechanism. This group has 12 elements having 4 irrudicible representations, with dimensions n_i, such that $\sum_i n_i = 12$. The characters of 4 representations are shown in table I. The complex number ω is the cube root of unity. Under A_4 fermions and scalars of this model [11] transform as $(\nu_i, l_i) \sim 3$, $l_i^c \sim 1, 1', 1''$, and $(\phi_i^0, \phi_i^-) \sim 3$. Charged lepton mass matrix in this model can be written as [12]:

TABLE I. Character table of A_4

Class	$\chi^{(1)}$	$\chi^{(2)}$	$\chi^{(3)}$	$\chi^{(4)}$
C_1	1	1	1	3
C_2	1	ω	ω^2	0
C_3	1	ω^2	ω	0
C_4	1	1	1	-1

$$M_l = \begin{pmatrix} h_1 v_1 & h_2 v_1 & h_3 v_1 \\ h_1 v_2 & h_2 v_2 & h_3 v_2 \\ h_1 v_3 & h_2 v_3 & h_3 v_3 \end{pmatrix}$$

$$= \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{pmatrix} \begin{pmatrix} h_1 & 0 & 0 \\ 0 & h_2 & 0 \\ 0 & 0 & h_3 \end{pmatrix} \sqrt{3} v, \tag{2}$$

for $v_1 = v_2 = v_3 = v$. Considering the charged lepton mass matrix for each allowed value, we have to consider the symmetry for M_ν. Let

$$M_\nu = \begin{pmatrix} a+b+c & f & e \\ f & a+b\omega+c\omega^2 & d \\ e & d & a+b\omega^2+c\omega \end{pmatrix} \tag{3}$$

be the Majorana neutrino mass matrix. Under A_4 a, b, c, and (d,e,f) comes from 1, $1'$, $1''$ and 3 respectively. Now using the case $b = c$ and $e = f = 0$ [13] and by an ansatz $a = 0$ and $d = 3b$ we can arrived to the expression of M_ν as mentioned by [10] is

$$M_\nu = m_{LL} = \begin{pmatrix} 2b & 0 & 0 \\ 0 & -b & 3b \\ 0 & 3b & -b \end{pmatrix}, \tag{4}$$

which gives the two-fold degenerate mass eigenvalues (m, m, m').

B. RGE for neutrino mass eigenvalues and mixing

The left-handed Majorana neutrino mass matrix m_{LL} which is generally obtained from seesaw mechanism at high scale M_R, is usually expressed in terms of $K(t)$, the coefficient of the dimension five neutrino mass operator [14, 15] in a scale-dependent manner, $t = \ln(\mu/1GeV)$ [16],

$$m_{LL}(t) = v_u^2 K(t), \tag{5}$$

where the vacuum expectation value (vev) is $v_u = v_0 \sin\beta$ and $v_0 = 174$ GeV in MSSM. The neutrino mass eigenvalues m_i and the PMNS mixing matrix U_{PMNS} [17] are then extracted through the diagonalisation of $m_{LL}(t)$ at every point in the energy scale t using equation (5),

$$m_{LL}^{diag} = Diag(m_1, m_2, m_3) = V_{\nu L} m_{LL} V_{\nu L}^T, \tag{6}$$

and the PMNS mixing matrix $U_{PMNS} = V_{\nu L}^\dagger$ in the basis where the charged lepton mass matrix is diagonal. The PMNS mixing matrix,

$$U_{PMNS} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix}, \tag{7}$$

is usually parametrised in terms of the product of three rotations $R(\theta_{23})$, $R(\theta_{13})$ and $R(\theta_{12})$, (neglecting CP violating phases) by

$$U_{PMNS} = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13} \\ -c_{23}s_{12} - c_{12}s_{13}s_{23} & c_{12}c_{23} - s_{12}s_{13}s_{23} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23} & -c_{12}s_{23} - c_{23}s_{13}s_{12} & c_{13}c_{23} \end{pmatrix}, \tag{8}$$

where $s_{ij} = \sin\theta_{ij}$ and $c_{ij} = \cos\theta_{ij}$ respectively.

The RGEs for the eigenvalues of coefficient $K(t)$ in equation (5), defined in the basis where the charged lepton mass matrix is diagonal, can be expressed as [18]

$$\frac{d}{dt}K_i = \frac{1}{16\pi^2}\sum_{f=e,\mu,\tau}[(-\frac{6}{5}g_1^2 - 6g_2^2 + 6Tr(h_u^2) + 2h_f^2U_{fi}^2]K_i. \tag{9}$$

Neglecting h_μ^2 and h_e^2 compared to h_τ^2, and taking scale-independent vev as in equation (5), we have the complete RGEs for three neutrino mass eigenvalues,

$$\frac{d}{dt}m_i = \frac{1}{16\pi^2}[(-\frac{6}{5}g_1^2 - 6g_2^2 + 6h_t^2) + 2h_\tau^2U_{\tau i}^2]m_i. \tag{10}$$

The above equations together with evolution equations for mixing angles (19-20), are used for numerical analysis in the present work.

The approximate analytical solution of equation (10) can be obtained by taking static mixing angle $U_{\tau i}^2$ in the integration range, as [19]

$$m_i(t_0) = m_i(t_R)exp(\frac{6}{5}I_{g1} + 6I_{g2} - 6I_t)exp(-2U_{\tau i}^2I_\tau). \tag{11}$$

The integrals in the above expression are usually defined as[16, 19]

$$I_{gi}(t_0) = \frac{1}{16\pi^2}\int_{t_0}^{t_R} g_i^2(t)dt, \tag{12}$$

and,

$$I_f(t_0) = \frac{1}{16\pi^2}\int_{t_0}^{t_R} h_f^2(t)dt, \tag{13}$$

where $i = 1, 2, 3$ and $f = t, b, \tau$ respectively. For a two-fold degenerate neutrino masses , viz., $m_{LL}^{diag} = Diag.(m, m, m') = U_{PMNS}^T m_{LL} U_{PMNS}$, equation (11) is further simplified to the following expressions

$$m_1(t_0) \approx m(t_R)(1 + 2\delta_\tau(c_{12}s_{13}c_{23} - s_{12}s_{23})^2) + O(\delta_\tau^2), \tag{14}$$

$$m_2(t_0) \approx -m(t_R)(1 + 2\delta_\tau(c_{23}s_{13}s_{12} + c_{12}s_{23})^2) + O(\delta_\tau^2), \tag{15}$$

$$m_3(t_0) \approx m'(t_R)(1 + 2\delta_\tau(c_{13}c_{23})^2) + O(\delta_\tau^2). \tag{16}$$

While deriving the above expressions, the following approximations are made,

$$\exp(-2|U_{\tau i}|^2I_\tau) \simeq 1 - 2|U_{\tau i}|^2I_\tau = 1 + 2|U_{\tau i}|^2\delta_\tau,$$

$$-\delta_\tau = I_\tau \simeq \frac{1}{\cos^2\beta}(m_\tau/4\pi v)^2 \ln(M_R/m_t),$$

The quantity δ_τ is always negative in MSSM and the approximation on δ_τ taken here is valid only if t_0 is associated with the top-quark mass. From equations (14) and (15), the low energy solar neutrino mass scale is then obtained as

$$\triangle\, m^2_{21}(t_0) = m^2_2 - m^2_1 \approx 4\delta_\tau m^2(\cos 2\theta_{12}(s^2_{23} - s^2_{13}c^2_{23}) + s_{13}\sin 2\theta_{12}\sin 2\theta_{23}) + O(\delta^2_\tau). \quad (17)$$

C. Evolution equations for mixing angles

The corresponding evolution equations for the PMNS matrix elements U_{fi} are given by [18]

$$\frac{dU_{fi}}{dt} = -\frac{1}{16\pi^2}\sum_{k\neq i}\frac{m_k+m_i}{m_k-m_i}U_{fk}(U^T H^2_e U)_{ki}, \quad (18)$$

where $f = e, \mu, \tau$ and $i, k = 1, 2, 3$ respectively. Here H_e is the Yukawa coupling matrices of the charged leptons in the diagonal basis and

$$(U^T H^2_e U)_{ki} = h^2_\tau(U^T_{k\tau}U_{\tau i}) + h^2_\mu(U^T_{k\mu}U_{\mu i}) + h^2_e(U^T_{ke}U_{ei}).$$

Neglecting h^2_μ and h^2_e as before, and denoting $A_{ki} = \frac{m_k+m_i}{m_k-m_i}$, equation (18) simplifies to [18]

$$\frac{ds_{12}}{dt} = \frac{1}{16\pi^2}h^2_\tau c_{12}[c_{23}s_{13}s_{12}U_{\tau 1}A_{31} - c_{23}s_{13}c_{12}U_{\tau 2}A_{32} + U_{\tau 1}U_{\tau 2}A_{21}], \quad (19)$$

$$\frac{ds_{13}}{dt} = \frac{1}{16\pi^2}h^2_\tau c_{23}c^2_{13}[c_{12}U_{\tau 1}A_{31} + s_{12}U_{\tau 2}A_{32}], \quad (20)$$

$$\frac{ds_{23}}{dt} = \frac{1}{16\pi^2}h^2_\tau c^2_{23}[-s_{12}U_{\tau 1}A_{31} + c_{12}U_{\tau 2}A_{32}]. \quad (21)$$

These equations are general and also independent of any neutrino mass model.

D. Specific model for two-fold degenerate case

We confine our analysis to a particular neutrino mass model[10] having two-fold degenerate neutrino mass eigenvalues , viz., $m_i = (m, m, m')$. Then the master equation in (20) can be further simplified for input values $m_3 = m' \neq 0$, $s_{13} = 0$,

$$\frac{ds_{13}}{dt} = \frac{1}{16\pi^2}h^2_\tau c_{23}c^2_{13}[c_{12}s_{12}s_{23}(A_{31} - A_{32})]. \quad (22)$$

In both cases the mixing angle s_{13} evolves with energy scale, and this in turn induces solar mass scale through equation (17).

III. NUMERICAL ANALYSIS AND RESULTS

For a complete numerical analysis of the RGEs, equations (10),(19)-(21) for neutrino masses and mixing angles, we follow here two consecutive steps (i) bottom-up running [19] in the first place, and then (ii) top-down running [16] in the next. In the first step (i), the running of the RGEs for the third family Yukawa couplings (h_t, h_b, h_τ) and three gauge couplings (g_1, g_2, g_3) in MSSM , are carried out from top-quark mass scale $(t_0 = \ln m_t)$ at low energy end to high energy scale M_R [19, 20]. In the present analysis we consider the high scale value as the unification scale $M_R = 1.6 \times 10^{16} GeV$, with different $\tan\beta$ input values to check the stability of the model at low energy scale. For simplicity of the calculation the SUSY breaking scale is taken at the top-quark mass scale $t_0 = \ln m_t$ [16, 19]. We adopt the standard procedure to get the values of gauge couplings at top-quark mass scale from the experimental CERN-LEP measurements at M_Z, using one-loop RGEs, assuming the existence of a one-light Higgs doublet and five quark flavours below m_t scale [19, 20].Using CERN-LEP data, $M_Z = 91.187 GeV$,$\alpha_s(M_Z) = 0.118 \pm 0.004$, $\alpha_1(M_Z) = 127.9 \pm 0.1$, $sin^2\theta_W(M_Z) = 0.2316 \pm 0.0003$, and SM relations,

$$\frac{1}{\alpha_1(M_Z)} = \frac{3}{5}\frac{(1 - sin^2\theta_W(M_Z))}{\alpha(M_Z)}, \frac{1}{\alpha_2(M_Z)} = \frac{sin^2\theta_W(M_Z)}{\alpha(M_Z)} \quad and, g_i^2 = 4\pi\alpha_i, \tag{23}$$

we calculate the gauge couplings at M_Z scale, $\alpha_1(M_Z) = 0.0169586$, $\alpha_2(M_Z) = 0.0337591$, $\alpha_3(M_Z) = 0.118$. As already mentioned, we consider the existence of one light Higgs doublet $(n_H = 1)$ and five quark flavours $(n_F = 5)$ in the scale $M_Z - m_t$. Using one-loop RGEs of gauge couplings, get $g_1(m_t) = 0.463751$, $g_2(m_t) = 0.6513289$ and $g_3(m_t) = 1.1891996$. Similarly, the Yukawa couplings are also evaluated at top-quark mass scale for input values of $m_t(m_t) = 174$ GeV, $m_b(m_b) = 4.25$ GeV, $m_t(m_\tau) = 1.785 GeV$ and the QED-QCD rescaling factors $\eta_b = 1.55$, $\eta_\tau = 1.015$ in the standard fashion [20],

$$h_t(m_t) = \frac{m_t(m_t)\sqrt{1 + tan^2\beta}}{174 tan\beta},$$

$$h_b(m_t) = \frac{m_b(m_t)\sqrt{1+tan^2\beta}}{174},$$
$$h_\tau(m_t) = \frac{m_\tau(m_t)\sqrt{1+tan^2\beta}}{174}. \tag{24}$$

where $m_b(m_t) = \frac{m_b(m_b)}{\eta_b}$, $m_\tau(m_t) = \frac{m_\tau(m_\tau)}{\eta_\tau}$. The one-loop renormalization group equations (RGEs) for top quark, bottom quark and τ-lepton Yukawa couplings in the MSSM in the range of mass scales $m_t \leq \mu \leq M_R$ are given by

$$\frac{d}{dt}h_t = \frac{h_t}{16\pi^2}(6h_t^2 + h_b^2 - \sum_{i=1}^{3} c_i g_i^2), \tag{25}$$

$$\frac{d}{dt}h_b = \frac{h_b}{16\pi^2}(6h_b^2 + h_\tau^2 + - \sum_{i=1}^{3} c_i^{'} g_i^2), \tag{26}$$

$$\frac{d}{dt}h_\tau = \frac{h_\tau}{16\pi^2}(4h_\tau^2 + 3h_b^2 - \sum_{i=1}^{3} c_i^{''} g_i^2), \tag{27}$$

where

$$c_i = \begin{pmatrix} \frac{13}{15} \\ 3 \\ \frac{16}{3} \end{pmatrix}, c_i^{'} = \begin{pmatrix} \frac{7}{15} \\ 3 \\ \frac{16}{3} \end{pmatrix} and, c_i^{''} = \begin{pmatrix} \frac{9}{5} \\ 3 \\ 0 \end{pmatrix}. \tag{28}$$

The two-loop RGEs for the gauge couplings are similarly expressed in the range of mass scales $m_t \leq \mu \leq M_R$ as

$$\frac{d}{dt}g_i = \frac{g_i}{16\pi^2}[b_i g_i^2 + \frac{1}{16\pi^2}(\sum_{j=1}^{3} b_{ij} g_i^2 g_j^2) - \sum_{j=t,b,\tau} a_{ij} g_i^2 h_j^2)], \tag{29}$$

where

$$b_i = \begin{pmatrix} 6.6 \\ 1 \\ -3 \end{pmatrix}, b_{ij} = \begin{pmatrix} 7.9 & 5.4 & 17 \\ 1.8 & 25 & 24 \\ 2.2 & 9 & 14 \end{pmatrix} and, a_{ij} = \begin{pmatrix} 5.2 & 2.8 & 3.6 \\ 6 & 6 & 2 \\ 4 & 4 & 0 \end{pmatrix}. \tag{30}$$

value of h_t, h_b, h_τ, g_1, g_2, g_3 evaluated at high scale $M_R = 1.6 \times 10^{16}$ from equation (25)-(27) and (29) are

$$h_t(M_R) = 0.977791667, h_b(M_R) = 1.03416789,$$
$$h_t(M_R) = 1.14931369, g_1(M_R) = 0.718215168,$$
$$g_t(M_R) = 0.726091146, g_3(M_R) = 0.725136459.$$

In the second step (ii), the runnings of three neutrino masses $(m_1, m_2, m_3)$and mixing angles (s_{12}, s_{23}, s_{13}) are carried out together with the running of Yukawa and gauge couplings, from high scale $t_R(= lnM_R)$ to low scale t_o. In this case, we use the input values of Yukawa and gauge couplings evaluated earlier at scale t_R from the first stage running of RGEs in case (i). In principle , one can evaluate neutrino masses and mixing angles at every point of the energy scale. The same analysis is repeated for different values of tanβ as well as the input value of high scale M_R. It can be noted that in the present problem, the running of other SUSY parameters such as M_0, $M_{1/2}$, μ, are not required and hence, it is not necessary to supply their input values. As emphasised earlier, we study here a two-fold degenerate neutrino mass model , $m_i = (m, m, m')$at high scale.

We are now interested here to study radiative generation of both $\triangle m^2_{21}$ and $\sin\theta_{13}$ for the case when $m' \neq 0$ and $s_{13} = 0$. Such studies can give the possible origin of the solar scale as well as the reactor angle with broken A_4 model. During the runnings of mass egenvalues and mixing angles in the RGEs in the present analysis, the non-zero input value of m' will first induce radiatively a non-zero values of s_{13} which in turn generates non-zero value of $\triangle m^2_{21}$ while running from high to low scale.

We compare our numerical results with recent neutrino oscillation data mentioned in section I and observed the following results:

- Variation of $sin^2\theta_{23}$ with energy shows that $sin^2\theta_{23}$ is stable at low energy for different tanβ values from 10 to 55 (figure 1).
- Variation of $sin^2\theta_{12}$ with energy shows that $sin^2\theta_{12}$ is stable at low energy for different tanβ values from 40 to 55 (figure 1).
- Variation of Δm^2_{23} with energy shows that Δm^2_{23} is stable at low energy for different tanβ values from 10 to 52 (figure 1).
- Generation of Δm^2_{21} with energy shows that Δm^2_{21} is stable at low energy for the tanβ values 50 to 55 (figure 1).
- Generation of $sin^2\theta_{13}$ with energy shows that $sin^2\theta_{13}$ is stable at low energy for the tanβ= 55 (figure 2).

It is observed from the figure that all the parametrs stable under radiative corrections with tan$\beta = 55$ except the fact that prediction for Δm^2_{23} are slightly deviated from 3σ range of



FIG. 1. Radiative generation of the $sin^2\theta_{12}$, $sin^2\theta_{23}$, Δm^2_{21} and Δm^2_{23} at low energy for different values of tanβ



FIG. 2. Radiative generation of $sin^2\theta_{13}$ for different tanβ values at low energy, and variation of $sin^2\theta_{13}$ with respect to $sin^2\theta_{12}$, $sin^2\theta_{23}$ and Δm^2_{21} for tanβ=55

experimental data. Therefore, it is very likely that more precise future data from neutrino oscillation experiments can expain these exceptions in the present analysis.

IV. DISCUSSION

Analysis of the effect of neutrino mass eigenvalues and mixings in case of two-fold degenerate neutrinos are done with RGE in the broken A_4 model. It is observed that all the parameters are stable at low slcale under radiative corrections for tanβ around 55 except the fact that for Δm^2_{23} the prediction are slightly below the 3σ range of experimental data. Apart from this exception, all other predicted values of the neutrino parameters are consistent with neutrino oscillation data. In view of above, the scenario of two-fold degenerate neutrinos can survive in nature within the framework of Type I seesaw mechanism. However, here we have not made any attempt to explain the CP violating phase. As an extension of this work, one can incorporate the RGE for phases which we have left for future studies.

[1] S. Fukuda et al. (Super-Kamiokande), Phys. Rev. Lett. **86**, 5656 (2001), hep-ex/0103033; Q. R. Ahmad et al. (SNO), Phys. Rev. Lett. **89**, 011301 (2002), nucl-ex/0204008; Phys. Rev. Lett. **89**, 011302 (2002), nucl-ex/0204009; J. N. Bahcall and C. Pena-Garay, New J. Phys. **6**, 63 (2004), hep-ph/0404061; K. Nakamura et al., J. Phys. **G37**, 075021 (2010).

[2] K. Abe et al. [T2K Collaboration], Phys. Rev. Lett. **107**, 041801 (2011). [arXiv:1106.2822 [hep-ex]].

[3] Y. Abe et al., arXiv:1112.6353 [hep-ex].

[4] F. P. An et al. [DAYA-BAY Collaboration], Phys. Rev. Lett. **108**, 171803 (2012) [arXiv:1203.1669 [hep-ex]].

[5] J. K. Ahn et al. [RENO Collaboration], Phys. Rev. Lett. **108**, 191802 (2012) [arXiv:1204.0626][hep-ex]].

[6] M. C. Gonzalez-Garcia, M. Maltoni, J. Salvado and T. Schwetz, JHEP **12**, 123 (2012)[arXiv:1209.3023 [hep-ph]].

[7] D. Borah, M. K. Das, Nucl. Phys. **B870** , 461 (2013);arXiv:1303.1758; M. K. Das, D. Borah, R. Mishra, Phys. Rev. **D86**, 095006 (2012).

[8] Y. BenTov, X. G. He, A. Zee, JHEP **12**, 093 (2012), arXiv:1208.1062; G. Altarelli, F. Feruglio, L. Merlo and E. Stamou, JHEP **1208**, 021 (2012); G. Altarelli and F. Feruglio, Phys. Rep. **320** (1999) 295; Y. Shimizu, M. Tanimoto, A. Watanabe, Prog. Theor. Phys. **126** 81 (2011).

[9] S. F. King, Phys. Lett. **718**, (1), 136 (2012); C. Duarah, A. Das, N. N. Singh, Jul 2012. 16 pp. Phys.Lett. **B718**, 147 (2012); N. K. Francis, N. N. Singh, Nucl. Phys. **B863**, 19 (2012); B. Brahmachari, A. Raychaudhuri, Phys.Rev. **D86**, 051302 (2012); S. Goswami, S. T. Petcov, S. Ray, W. Rodejohann, Phys.Rev. **D80**, 053013 (2009).

[10] see equation (28) of A. Zee, Phys. Lett. **B630**, 58 (2005).

[11] E. Ma, R. Rajasekaran, Phys. Rev. **D64**, 113012 (2001).

[12] E. Ma, Talk at VI-Silafac, Puerto Vallarta, November 2006, [arXiv:0612013].

[13] E. Ma, Phys. Rev. **D70**, 031901 (2004).

[14] P. Chankowski and Z. Pluciennik, Phys. Lett. **B316**, 312 (1993) ; K.S.Babu, C.N.Lung and J.Pantaleone, Phys.Lett. **B319**, 191 (1993). .

[15] S. Antusch, M. Drees, J. Kersten, M. Lindner and M. Ratz, Nucl. Phys. **B519**, 238 (2001) ; Phys. Lett. **B525**, 130 (2002).

[16] S. F. King and N. Nimai Singh, Nucl. Phys. **B591**, 3 (2000); Nucl. Phys. **B596**, 81 (2001).

[17] B. Pontecorvo, Sov. Phys. JETP **7**, 172 (1958); Z. Maki, M. Nakagawa, S. Sakata, Prog. Theor. Phys. **28**, 870 (1972).

[18] P. H. Chankowski, W. Krolikowski, S. Pokorski, Phys. Lett. **B473**, 109 (2000).

[19] M. K. Parida and N. Nimai Singh, Phys. Rev. **D59**, 032001 (1998).

[20] N. N. Singh, Eur. Phys. J. **C19**, 137 (2001).